

Pages (73-77)

CC BY NC

الهدف من هذه الدراسة هو التأكد من تركيز بعض المعادن الثقيلة في عينات الحليب المأخوذة من الماعز التي ترعى بالقرب من مدينة بني وليد ، ليبيا. بعد جمع عشر عينات من الحليب، تم أخذها على الفور إلى المختبر وتبريدها لمزيد من الفحص. باستخدام مطياف امتصاص الذهب (FAAS)، تم قياس تراكيزات الرصاص (Pb) والكاديوم (Cd) والنحاس (Cu) والحديد (Fe) في العينات. في حين كان الرصاص (Pb) والكاديوم (Cd) والنحاس (Cu) في جميع عينات الدراسة أقل من حدود الكشف الألي في جميع العينات، أظهرت النتائج أنه تم اكتشاف الحديد (Fe) في جميع العينات لآلآلآ، على لآلآ لآ لآ هذا المستولآ كان أقل من الحد الموصى به لآلآليآ من لآ لآ لآ لآلآلآ لآ لآلآلآ. (WHO) هذا بعد . أن حليب الماعز مناسب للاستهلاك.

of lactose, protein, fat, and total solids, there are notable differences in the size, content, and structure of casein micelles (Domagala, 2009). Special qualities of goat milk include its high digestibility, unique alkalinity, strong buffering capacity, and potential medical benefits (Ribeiro, & Ribeiro (2010), Getaneh (2016).goat milk contains lipids with more favorable physical properties than cow milk. The two primary ways that heavy metal pollution enters the human body are by ingestion and breathing (Tripathi *et al.*, 1997).Important elements

included in animal milk include phosphorus, calcium, potassium, magnesium, sodium, and chloride. Trace elements include iron, cadmium, nickel, copper, zinc, manganese, and manganese (Coni *et al.*, 1996). When high concentrations of heavy metals such as Cu, Cd, Zn, Ni, Hg, Pb, Fe As, and Cr are introduced to lactating animals, the metals accumulate in their milk and when consumed by humans, can have major health consequences (Onundi *et al.*, 2010). Microelements thought to be critical for human growth, such as copper (Cu), iron (Fe), zinc (Zn), and selenium (Se), are found in milk. However, if levels of heavy metals including arsenic (As), cadmium (Cd), and lead (Pb) exceed maximum allowable levels, it can negatively affect human health and welfare (Aslam *et al.*, 2011).

Due to the major impact on human health, it is now required to Detect and assess the amounts of heavy metals in milk as a result of rising environmental contamination (Caggiano *et al.*, 2005). Heavy metals have been found in milk and other food products, according to numerous investigations (Rahimi, 2013, Starska *et al* 2011, Ben-Hander *et al.*, 2020, Balli *et al.*, 2023). Determining trace elements is a difficult task because of the low concentrations of heavy metal ions and complex emulsions in milk.

Sources of heavy metal pollution can come from fertilizers polluted with cadmium, which is linked to the buildup of cadmium in animal offals, or through direct exposure to grass and crops raised for animal feed (Bramely, 1992). On the other hand, industrial pollution has been caused by contaminated water from factory waste discharge. Environmentally, as well, from the fumes released during mining or from cars running on leaded gasoline. According to Onundi *et al.* (2010), prolonged or excessive exposure to heavy metals can cause a variety of health problems, including cancer, behavioral issues, hepatotoxicity, neurotoxicity, vomiting, ulcerative colitis, Alzheimer's disease, and tissue damage and lung irritation.

Small levels of some metals, including lead, can have varying effects on people's health when they are regularly consumed. Since milk and dairy products are considered the primary sources of nutrition for children, it is crucial to monitor the level of trace elements in them. Complete removal or prevention of chemical contaminants from milk is not possible because the lipophilic contaminants will find their way into the persistent fat compounds from where heavy meta-bolic acids are found. For example, regular consumption of small amounts of lead may cause various impacts on the health of growing infants and children. These impacts include impaired concentration, retardation of mental development (such as reading and learning disabilities), adverse effects on kidney function, blood chemistry, and

cardiovascular system, and degradation of hearing. In addition, heavy metals accumulate in food chains through biotransformation, bioaccumulation, and biomagnifications because they are not biodegradable in the natural world (Aslam *et al.*, 2011). Thus, the purpose of this study was to ascertain the impact of certain heavy metals (Pb, Cd, Cu, and Fe) on public health.

MATERIALS AND METHODS

SAMPLES COLLECTION:

Ten milk samples from goats were randomly selected from various locations, particularly in the vicinity of Bani Waleed City in Libya. In order to be evaluated, Milk samples were collected in sterile polypropylene bottles of 50 ml then moved as quickly as possible in an ice box to the chemical laboratory at the Bani Waleed University Faculty of Education and they were frozen under -20°C until their analysis (Balli *et al.*.,2023).



Fig. 1 Bani Waleed City map shows sampling sites.(Google maps)

DETERMINATION OF SOME HEAVY METALS

PREPARATION OF SAMPLES:

2ml of milk was digested using a 1:3 ratio of H_2O_2 to HNO_3 . On a hot plate, the samples were heated until their volume dropped to two millilitres. The 2 ml sample solution was then turned into a clear solution by diluting it with 20 ml of deionized water. Using distilled water, the beaker's contents were increased to the necessary volume (25 ml), and a Flame Atomic Absorption Spectrophotometer was used to analyze the results (Richards, 1968).

RESULTS AND DISCUSSION

The physiological state of each lactation stage, as well as nutritional and environmental circumstances, can affect the mineral content of milk. Preliminary data suggests that consuming ovine and goat milk can provide valuable content in the heavy metals necessary for various metabolic activities. This is supported by comparisons with literature data related to these milk products. The obtained results in Table (1) illustrated that the minor essential element Fe was found in a range of (0.400-0.501ppm). Meanwhile, heavy metals (Pb, Cu, and Cd) could not be detected in examined goat milk because they were lower than instrumental detection limits in all samples.

Table (1): Results of some heavy metals (ppm) in goat milk.

Heavy metals	Min	Max	Mean $\pm$ SEM	WHO
lead	< 0.01	< 0.01	-	0.02
Cadmium	< 0.01	< 0.01	-	0.50
Copper	< 0.01	< 0.01	-	0.40
Iron	0.400	0.502	0.451 $\pm$ 0.01	0.50

0.01 = Detection limit

It appears from a more thorough examination of the data that the lead levels in the milk sample are lower than the experimental detection limits (Table 1). And those reported by other Authors since there are limited published data on harmful metal concentrations in goat milk. Lower Pb values of 7.37 μ g/L were previously recorded in Iran (Rahimi 2013) in comparison to the current results. However, exceptionally high Pb concentrations of 42.69 μ g/L were discovered in Pakistani industrial sites (Javed *et al.* 2009). The high Pb levels in the blood of goats grazing near highways or mining areas in Spain, which were 2.2-4.5 higher than in animals grazing in rural areas, illustrate the impact of Pb contamination in the environment (García-Fernández *et al.* 2003). Previous investigations on Pb contents in cow milk in Croatia found higher Pb values in the range of 23.06-70.56 μ g/kg (Sikiric *et al.* 2003). In addition, mean Pb concentrations in Croatia's northern and southern regions were substantially higher, at 58.7 and 36.2 μ g/L, respectively (Bilandžić *et al.* 2011). Low Pb concentrations in milk are almost definitely related to the fact that grazing pastures are located in rural locations, away from main road routes.

In terms of Cd, all samples exhibited levels of the metal that were below the limits of instrumental detection, showing that the Bani Waleed region is not at risk for toxicological consequences associated with Cd. While Licata *et al.* (2004) and Tripathi *et al.* (1999) reported

Cd levels in cow milk as 0.0228 and 0.00007 mg/L, respectively. Different researchers, like Baldini *et al.* (1990), Cerutti (1999), and Martino *et al.* (2001), observed Cd concentrations in cattle milk ranging from 0.0002 to 0.03 mg/L, which is significantly lower than the levels determined in this study, Hussain (2000) examined metal ion concentrations in goat milk and discovered that it was more polluted with Cd than cattle milk, 0.084 vs. 0.076 mg/L, respectively. The values of Cd in goat milk in the current investigation are supported by the values found by Coni *et al.* (1996) in goat milk (0.15 mg/L). Caggiano *et al.* (2005), on the other hand, detected 0.05 mg/L Cd residues in goat milk.

It is crucial to emphasize that, in terms of Cu values, all milk samples displayed concentrations that were lower than those discovered by other authors. Low levels of copper in milk samples are most likely caused by the zinc concentration of the feed, which messes with the animals' copper absorption mechanism (Baldini, *et al.*, 1990). These results are consistent with what other writers have reported (Baldini *et al.* 1990; Cerutti 1999).

As a necessary trace element, iron catalyzes a number of metabolic processes. Plays an important function in the transportation, storage, and consumption of oxygen as a component of hemoglobin, myoglobin, cytochromes, and other proteins. Moreover, it functions as a cofactor for several enzymes, and a shortage of it causes anemia (Meshref *et al.*, 2014). It was observed that the Fe (0.400-0.502 ppm) content in goat milk was lower than that found in some previous studies, Ahmad *et al.* (2017) found that the mean concentration of Fe in goat milk was higher at 0.95 $\pm$ 0.30 mg/kg compared to sheep milk's 0.59 $\pm$ 0.32 mg/kg. Conversely; Meshref *et al.* (2014) found that the range of Fe content in goat milk was (1.320–45.619).

Finally, the concentration of heavy metals (Pb, Cd, Cu, and Fe) in this study in all samples is still less than the permissible limit by WHO standards and LNCMS. (2016).

In any case, the levels of heavy metals discovered in goat milk samples from Bani Waleed suggest that the milk is safe for consumption, this is due to low/no pollution of these metals in these areas since Bani Waleed is a location with little industrial activity and, low levels of heavy metals are generally associated with this feature.

CONCLUSION

In conclusion, based on the findings of this study, it is possible to conclude that all tested goat milk samples contained low quantities of lead (Pb), cadmium (Cd),

and copper (Cu), which did not exceed the WHO-approved level and LNCSM.(2016). While iron (Fe) was detected in some samples, the quantities did not surpass the allowable limit. However, the number of heavy metals and samples evaluated in this study was restricted.

As a result, more research into "toxic" heavy metals in a larger number of fresh milk samples from different parts in Bani Waleed City is strongly advised.

REFERENCES

- Ahmad, I., Zaman, A., Samad, N., Ayaz, M. M., Rukh, S., Akbar, A., & Ullah, N. (2017). Atomic absorption spectrophotometry detection of heavy metals in milk of camel, cattle, buffalo and goat from various areas of Khyber-Pakhtunkhwa (KPK). *Pakistan. J Anal Bioanal Tech*, 8(3), 100367.
- Aslam, B., Javed, I., & Khan, F. H. (2011). Uptake of Heavy Metal Residues from Sewerage Sludge in the Milk of Goat and Cattle during Summer Season. *Pakistan Veterinary Journal*, 31(1).
- Baldini, M., Coni, E., Stacchini, A., & Stacchini, P. (1990). Presence and assessment of xenobiotic substances in milk and dairy products. *Annali Dell'istituto Superiore di Sanita*, 26(2), 167-175.
- Balli, N., Lebsir, D., Habila, S., Boutenoun, H., Boussof, L., Bouchehou, N., & Belkoucem, S. I. (2023). Health Risk Assessment of Heavy Metals in Goat's Milk for Adults Group in Jijel Province (Algeria). *Journal of Applied Biological Sciences*, 17(1), 93-110.
- Ben-Hander, G. M., Alsaeh, I. A., & Abdusalam, A. A. (2020). Primary study for determination of some toxic heavy metals in goat milk samples in Sirte, Libya. *J. Basic Appl. Chem*, 10, 1-5.
- Bilandžić, N., Đokić, M., Sedak, M., Solomun, B., Varenina, I., Knežević, Z., & Benić, M. (2011). Trace element levels in raw milk from northern and southern regions of Croatia. *Food chemistry*, 127(1), 63-66.
- Boyazoglu, J., Hatziminaoglou, I., & Morand-Fehr, P. (2005). The role of the goat in society: Past, present and perspectives for the future. *Small Ruminant Research*, 60(1-2), 13-23.
- Bramley, A. J. (1992). Milk hygiene and machine milking. *Machine milking and lactation.*, 373-398.
- Caggiano, R., Anastasio, A., Macchiato, M., Paino, S., Ragosta, M., & Sabia, S. (2005). Metal levels in milk, dairy products, fodder and organ samples collected in ovine farms of Southern Italy. *ENVIRONMENTAL RESEARCH*, 99(1), 48-57.
- Cerutti, G. (2006). *Residui, additivi e contaminanti degli alimenti*. Tecniche nuove.
- Coni, E., Bocca, A., Coppolelli, P., Caroli, S., Cavallucci, C., & Marinucci, M. T. (1996). Minor and trace element content in sheep and goat milk and dairy products. *Food Chemistry*, 57(2), 253-260.
- Domagała, J. (2009). Instrumental texture, syneresis and microstructure of yoghurts prepared from goat, cow and sheep milk. *International Journal of Food Properties*, 12(3), 605-615.
- Garcia-Fernandez, A. J., Navas, I., Romero, D., Gómez-Zapata, M., & Luna, A. (2003). Influence of leaded-gasoline regulations on the blood lead concentrations in Murciano-Granadina goats from Murcia Region, Southeast Spain. *Bulletin of environmental contamination and toxicology*, 70, 1178-1183.
- Getaneh, G., Mebrat, A., Wubie, A., & Kendie, H. (2016). Review on goat milk composition and its nutritive value. *Journal of Nutrition and Health Sciences*, 3(4), 1-10.
- Hussain, S. I. (2000). Irrigation of crops with sewage effluent: implications and movement of Pb and Cr as affected by soil texture, Lime, Gypsum and organic matter. *University of Agriculture Faisalabad Pakistan*.
- Javed, I., Jan, I., Muhammad, F., Zargham Khan, M., Aslam, B., & Sultan, J. I. (2009). Heavy metal residues in the milk of cattle and goats during winter season. *Bulletin of environmental contamination and toxicology*, 82, 616-620.
- Libyan National Center for Standardization and Metrology, Libyan Standard No. 357 /2016.
- Licata, P., Trombetta, D., Cristani, M., Giofre, F., Martino, D., Calo, M., & Naccari, F. (2004). Levels of "toxic" and "essential" metals in samples of bovine milk from various dairy farms in Calabria, Italy. *Environment international*, 30(1), 1-6.
- Martino, F. A. R., Sánchez, M. L. F., & Sanz-Medel, A. (2001). The potential of double focusing-ICP-MS for studying elemental distribution patterns in whole milk, skimmed milk and milk whey of

- different milks. *Analytica chimica acta*, 442(2), 191-200.
- Meshref, A. M., Moselhy, W. A., & Hassan, N. E. H. Y. (2014). Heavy metals and trace elements levels in milk and milk products. *Journal of food measurement and characterization*, 8, 381-388.
- Muehlhoff, E., & Bennett, A. (2013). *Milk and dairy products in human nutrition*.
- Onundi, Y. B., Mamun, A. A., Khatib, M. A., & Ahmed, Y. M. (2010). Adsorption of copper, nickel and lead ions from synthetic semiconductor industrial wastewater by palm shell activated carbon. *International Journal of Environmental Science & Technology*, 7, 751-758.
- Rahimi, E. (2013). Lead and cadmium concentrations in goat, cow, sheep, and buffalo milks from different regions of Iran. *Food chemistry*, 136(2), 389-391.
- Ribeiro, A. C., & Ribeiro, S. D. A. (2010). Specialty products made from goat milk. *Small Ruminant Research*, 89(2-3), 225-233.
- Richards LA (1968) Diagnosis and improvement of saline and alkaline soils, 1st edn. IBH Publications Company, New Delhi Agri. Handbook No. 60
- Sikirić, M., Brajenović, N., Pavlović, I., Havranek, J. L., & Plavljanić, N. (2003). Determination of metals in cow's milk by flame atomic absorption spectrophotometry. *Czech Journal of Animal Science*, 48(11), 481-486.
- Starska, K., Wojciechowska-Mazurek, M., Mania, M., Brulińska-Ostrowska, E., Biernat, U., & Karłowski, K. (2011). Noxious Elements in Milk and Milk Products in Poland. *Polish Journal of Environmental Studies*, 20(4).
- Tripathi, R. M., Raghunath, R., & Krishnamoorthy, T. M. (1997). Dietary intake of heavy metals in Bombay city, India. *Science of the Total Environment*, 208(3), 149-159.
- Tripathi, R. M., Raghunath, R., Sastry, V. N., & Krishnamoorthy, T. M. (1999). Daily intake of heavy metals by infants through milk and milk products. *Science of the total environment*, 227(2-3), 229-235